

Effects of culturing conditions on production of D-hydantoinase from recombinant *Escherichia coli*

Jan-Hsiung Huang¹, Wen-Hwei Hsu², Chun-Wei Chen¹

¹Institute of Microbiology and Biochemistry, National Taiwan University, Taipei; and ²Institute of Molecular Biology, National Chung-Hsing University, Taichung, Taiwan, ROC

Received: February 9, 2004 Revised: March 12, 2004 Accepted: April 26, 2004

The effects of culturing conditions on D-hydantoinase production by a recombinant *Escherichia coli* strain were investigated using a controlled fed-batch fermentation system. Glucose concentration and pH of the culture broth were maintained at less than 3.3 g/L and at 7.0, respectively, in a 5 L jar fermentor. The optimal composition of the batch medium was glucose, 0.25%; yeast extract, 0.75%; (NH₄)₂SO₄, 0.25%; KH₂PO₄, 0.2%. The optimal feeding solution was glucose, 60%; yeast extract, 30%; ammonia water, 9%. Following 25-h cultivation, 0.02 mM isopropyl-β-D-thiogalactopyranoside was added to induce *dht* gene expression and the temperature was shifted from 32 to 27°C to avoid inclusion body formation. The plasmid-harboring *dht* gene was found to be stably maintained in the *E. coli*. Under optimal conditions, a cell density of about 25 g dry cell weight/L and a high volumetric productivity of 8300 U/L/h could be achieved after 48 h culture at agitation and aeration rates of 1000 revolutions per minute and 1 vvm, respectively.

Key words: *Escherichia coli*, hydantoinase, inclusion bodies, plasmids, recombinant proteins

D-Hydantoinase (Dht) catalyzes the reversible stereoselective ring opening of dihydropyrimidine and those D,L-5-monosubstituted hydantoins structurally related to N-carbamoyl-D-amino acids [1]. These N-carbamoyl-D-amino acids can be converted into amino acids by reacting with nitrous acid [2] or transformation with N-carbamoyl-D-amino acid amidohydrolase [3] without changing the configuration. Optically active D-amino acids thus obtained are widely used in the pharmaceutical industry as intermediates for the synthesis of semi-synthetic antibiotics, peptide hormones, pyrethroids, pesticides, sweeteners, and therapeutic amino acids [1,4-7]. The enzyme is therefore of particular interest to researchers in the biotechnological industry. In recent years, there has been an increasing demand for D-phenylglycine or D-*p*-hydroxyphenylglycine for the production of semi-synthetic β-lactam antibiotics such as amoxicillin, aspoxicillin and cefbuperazone [8]. The β-lactams are by far the most widely used antibiotics and are effective against a variety of bacterial infections [9]. Over the past few decades, however, widespread resistant strains have evolved from most of the common

pathogenic species, thereby raising concerns about their threat to public health [10-12].

The precursor D-amino acids for these antibiotics are currently synthesized in a two-step process using Dht enzyme and D,L-5-monosubstituted hydantoins as the starting substrates [13,14]. The latter can be synthesized relatively inexpensively from aldehydes either by reaction with potassium cyanide and ammonium carbonate [15] or by a process developed by Yamada et al [16]. Dht activity has been detected in microorganisms, plants and animals [17]. However, microorganisms remain the most common sources of this enzyme [18]. At present, the industrial-scale production of Dht mainly utilizes bacterial species of genera such as *Agrobacterium*, *Arthrobacter*, *Bacillus*, *Corynebacterium*, *Comamonas*, *Pseudomonas*, *Serratia*, and *Streptomyces* [1,4,6,16,17,19-22]. The levels of enzyme in these wild-type strains are relatively low [23] and in general in the range 13,000-35,000 U/L [18,24]. A recombinant strain capable of producing a large amount of Dht is therefore highly desirable in order to reduce the cost of D-amino acids. Many efforts have been made to construct recombinant strains with high Dht activity [1,4,13,17,25]. However, the levels of enzyme vary from one strain to another and are in general between 5220 and 1,900,000 U/L [13,17,25].

Corresponding author: Dr. Jan-Hsiung Huang, Institute of Microbiology and Biochemistry, College of Life Science, National Taiwan University, Taipei, Taiwan 10617, ROC.
E-mail: jhhuang@ntu.edu.tw

A recombinant *Escherichia coli* strain producing a significant amount of Dht has been constructed [26]. This recombinant *E. coli* strain harbors the *dht* gene from *Agrobacterium radiobacter* under the control of the T5 promoter. In this study, we investigated the optimal culture conditions for Dht production in shaker flasks or a benchtop fermentor under both batch and fed-batch operations. Furthermore, we report a fed-batch procedure that allows high cell density culture and leads to high-level expression of Dht.

Materials and Methods

Chemicals

5-(4-Hydroxyphenyl)-hydantoin (HPH) was obtained from Tokyo Chemical Industry (Tokyo, Japan). The N-carbamoyl-D-*p*-hydroxyphenylglycine (NC-D-*p*-HPG) standard was synthesized and kindly provided by Dr. Wu-Yung Yang (Union Chemical Laboratories, Industrial Technology Research Institute, Hsinchu, Taiwan). Yeast extract, tryptone and agar were obtained from Difco Laboratories (Detroit, MI, USA). Hydantoin, Tris, ampicillin (sodium salt), tetracycline (free base), isopropyl β -D-thiogalactoside (IPTG), bromthymol blue (sodium salt; BTB) and Coomassie brilliant blue R-250 were purchased from Sigma Chemical Co. (St Louis, MO, USA). Acrylamide was obtained from Merck KGaA (Darmstadt, Germany). All other chemicals were of reagent grade and were purchased from commercial suppliers.

Bacterial strain

A 1.4-kb *Kpn* 1-*Pst* 1 fragment containing the *dht* gene of *Agrobacterium radiobacter* DUL-DH 101 and the sequence encoding the N-terminal His-tag was cloned into expression vector pQE30 (Qiagen, Valencia, CA, USA) to generate pQE30/*dht* using standard procedures [27]. NovaBlue *E. coli* (*endA1 hsdR17* ($r_{k12}^- m_{k12}^-$) *supE44 thi-1 recA1 gyrA96 relA1 lac F'* [*proA*⁺*B*⁺ *lacI*^q *lacZ* Δ M15::Tn10(Tet^r)]), Novagen Inc. (Madison, WI, USA) was transformed with plasmid pQE30/*dht* using a classical CaCl₂ preparation [26,27]. The *dht* gene transcription is under the control of the phage T5 promoter and *lac* operator (Qiagen) and can be induced with IPTG. The transformant, referred to as *E. coli* NovaBlue(pQE30/*dht*) was used throughout this study. The strain was maintained as 40% (v/v) glycerol stock at -30°C in Luria-Bertani (LB) medium and was very stable throughout the whole experiment.

Media

LB medium [27] was employed for the propagation of *E. coli* transformants under the uninduced state. When necessary, ampicillin and tetracycline were added to the medium at concentrations of 100 and 10 μ g/mL, respectively. Modified LB (MLB) medium was LB medium supplemented with the following per liter: 10 g glucose, 0.1 g ampicillin and 0.01 g tetracycline. Semi-synthetic production (SSP) medium contained the following per liter: 10 g glucose, 10 g yeast extract, 7 g K₂HPO₄, 3 g KH₂PO₄, 0.1 g ampicillin and 0.01 g tetracycline. A synthetic medium modified from that of Horn et al [28] contained the following per liter: 5 g glucose, 0.5 g NaCl, 2 g (NH₄)₂SO₄, 9 g Na₂HPO₄, 3 g K₂HPO₄ and trace elements of 60 mg/L Fe(III) citrate, 3 mg/L H₃BO₃, 15 mg/L MnCl₂·4H₂O, 8.4 mg/L EDTA·2Na, 2 mg/L CuSO₄·5H₂O, 2.5 mg/L Na₂MoO₄·2H₂O, 2.5 mg/L CoCl₂·6H₂O, 8 mg/L ZnCl₂. Sterilization was performed at 121°C for 20 min. Glucose was autoclaved separately from the rest of the medium and added into the medium prior to inoculation. Ampicillin and tetracycline were sterilized by filtration.

Culture conditions

For cell activation, 0.2 mL frozen stock culture was added to 25 mL of MLB medium in a 50 mL Falcon tube (Falcon Plastics, Oxnard, CA, USA) and the tube shaken at 37°C and 125 rpm for 12 h. The inoculum culture was then prepared by adding 0.1% (v/v) of this reactivated culture to 50 mL MLB medium in a 250 mL Erlenmeyer flask, followed by culturing under the same conditions for 12 h. For shaker-flask culture, 2 mL of inoculum culture was added to 100 mL production medium in a 500-mL Hinton flask. The cultures were incubated at 37°C in a rotary shaker set at 125 rpm and 7 cm stroke (Model 707R, Hotech Instruments Corp., Taipei, Taiwan). After 6 h cultivation, 0.02 mM IPTG was added and the flasks were moved to another shaker at 27°C and 140 rpm (Model 706R, Hotech Instruments Corp., stroke 3.85 cm) for enzyme induction. For large-scale experiments, batch and fed-batch fermentations were performed in a classical 5 L jar fermentor (Model LS-205, Hotech Instruments Corp.) with a digital measurement and control unit from Suntex Instruments Co., Ltd. (Taipei, Taiwan). Antifoam 289 (Sigma A-5551) was automatically added as necessary to control foaming in the fermentor. Fifty mL of inoculum culture was added to 2 L of initial fermentation medium and the cultivation was carried

out under the conditions used for the shaker-flask culture. The aeration was kept constant at 2.0 L air/min, and agitation at 1000 rpm. The pH-stat method was employed for the control of substrate feeding. Medium pH was adjusted to 6.9-7.1 during the cultivation processes by the addition of either 9% (v/v) ammonia water or solutions of 600 g/L glucose and 300 g/L yeast extract with a peristaltic pump. The pH of the culture increased as glucose was depleted. When the pH exceeded the set high limit of 7.1, an equal volume of glucose solution and yeast extract solution was automatically added so that the glucose concentration in the culture broth was increased while the pH was decreased. In the fed-batch culture, the concentration of glucose in the medium was maintained below 3.3 g/L. When the pH was decreased below the lower limit of 6.9, ammonia water was automatically added to recover pH as well as supply a nitrogen source. No nutrient was fed when the pH was between 6.9 and 7.1. Antibiotics were not used in the fermentor study. The dissolved oxygen level was determined with a steam-sterilizable polarographic electrode (InPro 6000 series O₂ sensors; Mettler-Toledo GmbH, Switzerland), and the pH of culture broth was measured with a steam-sterilizable pH probe (InPro 3030/325 combination pH electrode; Mettler-Toledo GmbH, Switzerland). The time points of feeding and harvesting are reported in the Results section. Expression of the *dht* gene was induced by adding IPTG to a final concentration of 0.02 mM or as indicated. After induction with IPTG, cells were cultured for an additional 30 or more hours and harvested by centrifugation. Throughout the cultivation, pH, temperature, and dissolved oxygen were continuously monitored and recorded. Samples were automatically taken at various intervals for analysis.

Preparation of crude extracts by ultrasonication

The cells were obtained by centrifugation (2000 × g, 15 min), washed twice with 0.05 M Tris-HCl buffer (pH 8.0), then resuspended in the same buffer. Aliquots of 2 mL cell suspension corresponding to 30 absorbance [optical density (OD₆₀₀)] units were then disrupted with a cell sonicator (Model XL2020, 419A microtip probe; Misonix Inc., NY, USA) for 2 sec every 3 sec for a total of 165 sec in an ice bath. After centrifugation at 12,000 × g for 5 min at 4°C, the supernatants (soluble protein fraction) were used as the crude enzyme preparations. The pellets (inclusion body fraction) were washed twice with buffer and

resuspended in an equal volume of the buffer. The supernatants and pellets were analyzed by sodium dodecyl sulfate-polyacrylamide gel (SDS-PAGE) [29]. The enzyme with a 6-histidine-residue tag at the amino-terminal end was further purified from the supernatants on Ni-affinity columns as described by Hsu et al [30].

Assay of D-hydantoinase activity

Fifty μL of diluted supernatant or partially purified enzyme solution was incubated with 1% HPH (molecular weight [MW] 192.17) in 0.05 M Tris-HCl buffer (pH 8.0) in a total volume of 1 mL. The reaction was allowed to proceed at 37°C for 10 min in a reciprocating shaker water bath at 100 rpm, 3.85 cm stroke, and was terminated and deproteinized by the addition of 200 μL of 12% trichloroacetic acid (TCA). After removing the precipitate by centrifugation (8500 × g, 5 min), the amount of NC-D-*p*-HPG (MW 210.16) in the supernatant was quantified. One unit of Dht activity was defined as the amount of enzyme required to produce 1 μmole of NC-D-*p*-HPG per min under the assay conditions. A blank with TCA-denatured enzyme was assayed for all samples. Dht activity in colonies was detected on an LBHBTB agar plate, a blue plate containing LB supplemented with 10.0 g/L hydantoin, 0.1 g/L BTB, 1.0 mL 1 M MnCl₂·4H₂O, 0.25 mL 20 mM IPTG, 1.0 mL 1% tetracycline and 20.0 g/L Bacto-agar. Colonies that express Dht convert hydantoin to hydantoic acid, thus lowering the pH, and a yellowish zone was observed in the plate around the Dht-positive colonies against a blue background after 36 h incubation at 32°C.

Analytical methods

Dry cell weight (DCW) was measured after the cells were washed twice with distilled water and dried to a constant weight at 105°C. Cell density was monitored by measuring the optical density of the culture broth at 600 nm (OD₆₀₀) using a spectrophotometer. The culture broth was generally diluted with distilled water to yield an OD₆₀₀ value of 0.2-0.5. Biomass was calculated from the OD₆₀₀ and converted to DCW per liter according to predetermined calibration data. One OD unit corresponds to 0.33 ± 0.05 gDCW/L. The titers of total viable and ampicillin-resistant cells were determined by plating serial 10-fold diluted samples on LB agar plates in the presence or absence of 100 μg/mL ampicillin in triplicate. The number of colony-forming units was determined after incubating the plates at 37°C for 18 h. Glucose concentrations were measured

colorimetrically using dinitrosalicylic acid reagent [31]. Protein concentration was measured by the method of Bradford [32], using bovine serum albumin as a standard. SDS-PAGE was performed according to the method of Laemmli [29], using 12.5% polyacrylamide gels. Proteins were visualized by staining with Coomassie brilliant blue R-250 (Sigma) and quantified by densitometric analysis of stained bands on the gels with a Kodak Digital Science Electrophoresis Documentation and Analysis System.

NC-D-*p*-HPG was quantitated by high-performance liquid chromatography (HPLC) or estimated spectrophotometrically at 450 nm using Ehrlich's reagent (10% *p*-dimethylaminobenzaldehyde in 6.0 N HCl) as described by Grifantini et al [13]. HPLC analysis was performed on a reverse phase C18 column (4.6 × 250 mm; Merck, USA) [33]. Samples were eluted isocratically with acetonitrile:water (95:5 v/v) at a flow rate of 0.8 mL/min and detected at 274 nm using an ABI 783A UV detector (Applied Biosystems, Inc., Foster City, CA, USA). The detector was coupled to a chromatocorder for quantification. The retention times for NC-D-*p*-HPG and HPH were 6.5 and 9.8 min, respectively. The correlation coefficient of concentration and peak area for NC-D-*p*-HPG was 0.9997. The results obtained by colorimetric method and HPLC were in good agreement, with the variation around 3%.

Stability of the recombinant plasmid was determined by culturing plasmid-bearing *E. coli* in a 500 mL Hinton flask containing 100 mL SSP in the absence of ampicillin. Cells (0.1 mL) were inoculated into 100 mL of the next sterile medium. The proportion of Dht-producing cells in the population was determined by plating the cells on LBHBTB agar plates and incubating at 32°C for 36 h.

Results

Plasmid stability

The plasmid stability of *E. coli* NovaBlue(pQE30/*dht*) was investigated in SSP medium in the absence of ampicillin. At various intervals after inoculation, the proportions of Dht-producing cells in serially diluted samples were determined by plating the cells on LBHBTB agar plates. The development of yellowish color around each colony indicated that there was no loss in its capability to produce Dht. Plate assay indicated that the recombinant plasmid was stable over 200 generations in the ampicillin-free SSP medium. All colonies maintained the Dht plasmid after 200

generations. In addition, Dht levels were comparable among the different subcultures.

Effects of medium composition on cell growth

LB medium, a synthetic medium and SSP medium were used to evaluate the influence of medium composition on cell growth in 500 mL Hinton flasks. Although there was little or no growth of *E. coli* NovaBlue(pQE30/*dht*) in the synthetic medium because the strain used requires thiamine, good growth rates and high cell density were achieved in both LB and SSP media. The SSP medium was selected because of its lower cost and simple composition. The presence of yeast extract significantly increased the growth rate of this culture. With increasing amounts of yeast extract, the cell concentration and the specific growth rate increased, but the growth yields decreased in terms of yeast extract concentration (Table 1). The cell mass increased with increasing concentrations of yeast extract up to 7.5 g/L. Other components were tested but no significant effects were observed.

IPTG induction

Shake flask experiments were conducted to evaluate the influence of IPTG on the induction of Dht production in *E. coli* NovaBlue(pQE30/*dht*). The optimal temperature and IPTG concentration for Dht expression were found to be 27°C and 0.02 mM, respectively (Fig. 1). Low-level induction by IPTG allowed the growth of the recombinant strain to continue, thus prolonging the production phase of the induced cells. Lower temperature reduced the growth rate, therefore minimizing the formation of inhibitory byproducts such as acetate, and reducing the demand for oxygen. Even in the absence of IPTG, some amount of Dht was expressed at 27°C or 37°C. In addition, the formation of inclusion bodies was also significantly reduced (Fig. 2). Fig. 3 further shows that the *dht* gene could be efficiently induced with 0.02 mM IPTG in the mid-exponential phase of cell growth (at around

Table 1. Effects of yeast extract concentration on the growth rate of recombinant *Escherichia coli* NovaBlue(pQE30/*dht*)

Yeast extract (g/L)	μ (h ⁻¹)	Optical density (OD ₆₀₀)	Growth yield
0	0.267	0.29	-
2.5	0.611	2.40	0.96
5.0	0.675	3.26	0.65
7.5	0.776	4.51	0.60
10	0.805	4.80	0.48




Fig. 1. Effects of isopropyl β -D-thiogalactoside (IPTG) concentration and induction temperature on D-hydantoinase (Dht) expression. *E. coli* NovaBlue(pQE30/*dht*) was grown in Luria-Bertani medium in shaker flasks at 37°C under 125 rpm agitation. Following 6-h cultivation, different amounts of IPTG were added. The flask, containing two-thirds of the original culture, was transferred to another shaker set at 27°C and 140 rpm and cultured for an additional 6 h. The Dht activity of the culture broth was determined as described in Materials and Methods. The highest activity was taken as 100%. The percent activities under various IPTG concentrations and temperature conditions are shown. The mean value of at least 2 experiments is indicated by column height. The variation was less than 10%.

6 h). The maximal productivity, obtained when induced at 6 h following inoculation, was about 2.6-fold that obtained when IPTG induction was carried out 3 h earlier in the log phase of growth. No Dht activity was detected when cells were induced at the late log phase of growth (at 9 h).

D-Hydantoinase production

Based on the results described above, optimal conditions for Dht production of *E. coli* NovaBlue(pQE30/*dht*) in a 5 L jar fermentor were as follows. The optimal composition of batch medium was glucose, 0.25%; yeast extract, 0.75%; $(\text{NH}_4)_2\text{SO}_4$, 0.25%; KH_2PO_4 , 0.2%, while that for the feeding solution was glucose, 60%; yeast extract, 30%; ammonia water, 9%. During cultivation, the agitation and aeration rates were kept at 1000 rpm and 1 vvm, respectively and the pH was controlled automatically at 7.0 ± 0.1 , as described in Materials and Methods. Following 25-h cultivation, which was about the mid-log growth phase, 0.02 mM IPTG was added the temperature was shifted from 32 to 27°C. Under these conditions, a typical time course of the cultivation of *E. coli* NovaBlue(pQE30/*dht*) is shown




Fig. 2. Sodium dodecyl sulfate-polyacrylamide gel (SDS-PAGE) analysis of the soluble (A) and insoluble (B) crude extracts of *E. coli* NovaBlue(pQE30/*dht*) induced at different temperatures. The positions of molecular size markers in kDa are indicated on the right. Cells were grown in semi-synthetic production medium in shake flasks at 37°C under 125 rpm agitation. Following 6-h cultivation, 0.02 mM isopropyl β -D-thiogalactoside was added, and the flask was transferred immediately to another shaker, set at the different temperatures shown below, for enzyme induction. The cells were then harvested and lysed as described in Materials and Methods. Ten μg of protein was loaded in each lane on a 12.5% SDS-PAGE gel as described. Lane 1A and 1B, 27°C; lane 2A and 2B, 30°C; lane 3A and 3B, 32°C; lane 4A and 4B, 37°C; lane M, molecular mass markers in kDa. Bands corresponding to the 60 kDa protein are indicated by the arrow.

in Fig. 4. The specific growth rate was determined from the slope of the semi-logarithmic plot of cell density (OD_{600}) versus cultivation time. During the first 10 h of culture, a high specific growth rate of 0.423/h was obtained. However, the specific growth rate decreased to 0.026/h due to a combination of factors including IPTG induction, down-shift of temperature, and presumably carbon, oxygen, and growth factor depletion. The beginning of this second phase coincided with the onset of initial glucose depletion. Glucose was fed at this stage to maintain the maximum cell concentration for intracellular protein synthesis. To avoid excessive feeding of glucose, which is known to cause acetate accumulation [34], the nutrient feeding solution was added only when the glucose concentration declined to a certain level as indicated by an increase in pH. The highest Dht activity of about 400,000 U/L or 5 g/L of Dht was attained within 48 h and the activity remained relatively high even at 54 h. The maximum cell concentration reached around OD_{600} 75 (39 h) or 25 gDCW/L. A summary of Dht production under various culture conditions in *E. coli* NovaBlue(pQE30/*dht*) is shown in Table 2.




Fig. 3. Effect of induction time on D-hydantoinase (Dht) expression. Cells were grown in semi-synthetic production medium at 37°C at an agitation rate of 125 rpm. After 3- (●), 6- (○), or 9-h (△) cultivation, 0.02 mM isopropyl β-D-thiogalactoside was added, and the flask was transferred immediately to another shaker at conditions of 27°C and 140 rpm, for enzyme induction. At different time intervals, the Dht activity of the culture broth was determined as described in Materials and Methods. The highest activity was taken as 100%.

SDS-PAGE analysis of Dht

In order to determine if all Dht was appropriately processed to an active state, aliquots of cultures under different induction temperatures were analyzed by SDS-PAGE (Fig. 2). A prominent protein band with an apparent molecular mass of 60 kDa, corresponding to

the subunit MW of Dht was observed [35]. The 6-His-tagged-Dht was purified to homogeneity by His-Bind resin affinity chromatography. Purified Dht had D-hydantoinase activity also showed a single band with an apparent molecular mass of approximately 60 kDa. After 6 h induction at 27°C, about 80% of the enzyme




Fig. 4. Profile of D-hydantoinase (Dht) production by a fed-batch high cell density culture of *Escherichia coli* NovaBlue(pQE30/*dht*). The composition (per L) of the initial fermentation medium was: 2.5 g glucose, 7.5 g yeast extract, 2.5 g (NH₄)₂SO₄, 2.0 g KH₂PO₄; and that of the feeding solution was: 600 g glucose, 300 g yeast extract, 333 mL of 28% ammonia water. The time courses of biomass [optical density (OD₆₀₀, ○)], glucose (—), Dht activity (▲) as well as specific growth rates are shown. The closed box symbol (□) represents log OD₆₀₀, which was used to determine the specific growth rate. Agitation speed and aeration rate were kept constant at 1000 rpm and 1 vvm, respectively, throughout the cultivation. Glucose concentration and pH were controlled at below 3.3 g/L and 6.9-7.1, respectively. The time of isopropyl β-D-thiogalactoside addition, followed immediately by temperature shift from 32 to 27°C, is indicated by the arrow.

Table 2. Summary of D-hydantoinase production under different culture conditions in *Escherichia coli* NovaBlue(pQE30/dht)

	Hinton flask		5 L fermentor
	Luria-Bertani medium	Semi-synthetic production medium	High cell density culture
Isopropyl β -D-thiogalactoside (mM)	0.02	0.02	0.02
Cultivation time (h)	32	32	48
Final optical density (OD ₆₀₀)	7.3	12	75
Yield (U/g glucose)	-	6500	4950
Specific activity (U/g dry cell weight)	6100	16,400	16,200
Productivity (U/L/h)	460	2000	8300
D-hydantoinase activity (U/L)	14,700	65,000	400,000
Increase of Dht activity	-	4.42-fold	27.2-fold
Yield of purified enzyme (g/L)	ND	ND	5

Abbreviation: ND = not determined

was present as active and soluble protein (Fig. 2, lane 1A and 1B). The ratios of soluble protein to total Dht were 63% (Fig. 2, lane 2A and 2B) and 51% (Fig. 2, lane 3A and 3B) when induction temperatures were 30°C and 32°C, respectively. No soluble Dht protein was detectable when the induction temperature was 37°C (Fig. 2, lane 4A and 4B).

Discussion

Hydantoinases belong to the family of cyclic amidases. They are closely related to dihydropyrimidinases and most likely take part in the catabolism of pyrimidines and thus might be harmful to cells. In this study, the optimal culture conditions to allow the production of a significant amount of the active and soluble form of Dht were investigated. The application of these modes of fed-batch cultivation resulted in a 6 to 7-fold increase in Dht activity compared to shake flask cultivation (Table 2). The enzyme production process can be divided into 2 phases (Fig. 4). A rapid enzyme production phase positively correlated to the late exponential phase of growth, which was followed by a slower phase of production resulting from a slower growth rate. No significant toxicity of the over-expressed foreign protein to the host was observed. Dht activity continued to increase even after the stationary phase was reached. After the maximal cell concentration was reached, the culture was maintained for an additional 20-25 h. During this period, no loss in enzyme activity or significant lysis was observed, as the cell concentration remained relatively constant at 25.0 ± 0.2 g/L and the activity of Dht in the culture broth was about 0.2% of the total. Under simple culture conditions, at a constant feed rate without oxygen enrichment, a satisfactory quantity of

total Dht (5 g/L) could be obtained. Dht activity as high as 1,900,000 U/L has been reported from a recombinant *E. coli* [25]. However, the enzymatic activity in that report was defined as the hydrolysis of hydantoin, instead of that of D,L-*p*-hydroxyphenylhydantoin which is utilized in industry and in our study. The advantageous features of *E. coli* NovaBlue(pQE30/dht) for industrial scale enzyme production include its ability to stably retain plasmid and to produce significant amounts of Dht from an antibiotic-free medium. Our results also showed that not only was the yield of the recombinant product high but also the fraction of insoluble, functionally inactive, protein was generally quite low (Fig. 2). The production of industrial enzymes as an insoluble form is obviously undesirable, as additional (and generally inefficient) measures are required to refold the protein. Furthermore, the enzyme could be easily purified by a 1-step affinity chromatography. The crude or purified overproduced Dht can be further utilized, either in solution or in immobilized form, for the production of D-amino acids.

Acknowledgment

This study was supported by a grant NSC 87-2316-B002-006-BI from the National Science Council, Taiwan, ROC.

References

- Syldatk C, May O, Altenbuchner J, Mattes R, Siemann M. Microbial hydantoinases – industrial enzymes from the origin of life? *Appl Microbiol Biotechnol* 1999;51:293-309.
- Takahashi S, Ohashi T, Kii Y, Kumagai H, Yamada H. Microbial transformation of hydantoins to N-carbamyl-D-amino acids. *J Ferment Technol* 1979;57:328-32.

3. Olivieri R, Fascetti E, Angelini L, Degen L. Microbial transformation of racemic hydantoins to D-amino acids. *Biotechnol Bioeng* 1981;23:2173-83.
4. Ogawa J, Shimizu S. Stereoselective synthesis using hydantoinases and carbamoylases. In: Patel RN, ed. *Stereoselective biocatalysis*. New York: Marcel Dekker; 2000:1-21.
5. Soong CL, Ogawa J, Honda M, Shimizu S. Cyclic-imide-hydrolyzing activity of D-hydantoinase from *Blastobacter* sp. Strain A17p-4. *Appl Environ Microbiol* 1999;65:1459-62.
6. Sylđatk C, Laufer A, Muller R, Hoke H. Production of optically pure D- and L- α -amino acids by bioconversion of D,L-5-monosubstituted hydantoin derivatives. *Adv Biochem Eng Biotechnol* 1990;41:29-75.
7. Yagasaki M, Ozaki A. Industrial biotransformations for the production of D-amino acids. *J Mol Catal, B Enzym* 1998; 4:1-11.
8. Sudge SS, Bastawde KB, Gokhale DV, Kalkote UR, Ravindranathan T. Production of D-hydantoinase by halophilic *Pseudomonas* sp. NCIM 5109. *Appl Microbiol Biotechnol* 1998;49:594-9.
9. Demain AL, Elander RP. The β -lactam antibiotics: past, present, and future. *Antonie van Leeuwenhoek* 1999;75:5-19.
10. Bradford PA. Extended-spectrum β -lactamases in the 21st century: characterization, epidemiology, and detection of this important resistance threat. *Clin Microbiol Rev* 2001;14: 933-51.
11. Cohen ML. Epidemiology of drug resistance: Implications for a post-antimicrobial era. *Science* 1992;257:1050-5.
12. Gold HS, Moellering RC Jr. Antimicrobial-drug resistance. *N Engl J Med* 1996;335:1445-53.
13. Grifantini R, Galli G, Carpani G, Pratesi C, Frascotti G, Grandi G. Efficient conversion of 5-substituted hydantoins to D- α -amino acids using recombinant *Escherichia coli* strains. *Microbiology* 1998;144:947-54.
14. Ogawa J, Shimizu S. Diversity and versatility of microbial hydantoin-transforming enzymes. *J Mol Catal, B Enzym* 1997; 2:163-76.
15. Morin A, Hummel W, Kula MR. Production of hydantoinase from *Pseudomonas fluorescens* strain DSM 84. *Appl Microbiol Biotechnol* 1986;25:91-6.
16. Yamada H, Takahashi S, Kii Y, Kumagai H. Distribution of hydantoin hydrolyzing activity in microorganisms. *J Ferment Technol* 1978;56:484-91.
17. LaPointe G, Viau S, Leblanc D, Robert N, Morin A. Cloning, sequencing, and expression in *Escherichia coli* of the D-hydantoinase gene from *Pseudomonas putida* and distribution of homologous genes in other microorganisms. *Appl Environ Microbiol* 1994;60:888-95.
18. Morin A, Tran Trung NH, LaPointe G, Dubeau H. Conditions used with a continuous cultivation system to screen for D-hydantoinase-producing microorganisms. *Appl Microbiol Biotechnol* 1995;43:259-66.
19. Matsuda K, Sakata S, Kaneko M, Hamajima N, Nonaka M, Sasaki M, et al. Molecular cloning and sequencing of a cDNA encoding dihydropyrimidinase from the rat liver. *Biochim Biophys Acta* 1996;1307:140-4.
20. May O, Nguyen PT, Arnold FH. Inverting enantioselectivity by directed evolution of hydantoinase for improved production of L-methionine. *Nature Biotechnol* 2000;18:317-20.
21. Morin A, Leblanc D, Paleczek A, Hummel W, Kula MR. Comparison of seven microbial D-hydantoinases. *J Biotechnol* 1990;16:37-48.
22. Takahashi S, Kii Y, Kumagai H, Yamada H. Purification, crystallization and properties of hydantoinase from *Pseudomonas striata*. *J Ferment Technol* 1978;56:492-8.
23. Hartley CJ, Manford F, Burton SG, Dorrington RA. Overproduction of hydantoinase and N-carbamoylamino acid amidohydrolase enzymes by regulatory mutants of *Agrobacterium tumefaciens*. *Appl Microbiol Biotechnol* 2001; 57:43-9.
24. Achary A, Hariharan KA, Bandhyopadhyaya S, Ramachandran R, Jayaraman K. Application of numerical modeling for the development of optimized complex medium for D-hydantoinase production from *Agrobacterium radiobacter* NRRL B 11291. *Biotechnol Bioeng* 1997;55:148-54.
25. Lee DC, Kim GJ, Cha YK, Lee CY, Kim HS. Mass production of thermostable D-hydantoinase by batch culture of recombinant *Escherichia coli* with a constitutive expression system. *Biotechnol Bioeng* 1997;56:449-55.
26. Perng RY. Characterization and protein engineering of D-hydantoinase from *Agrobacterium radiobacter* and mutational analysis of the catalytic sites of D-hydantoinase from *Bacillus caldolyticus*. M.S. thesis, National Chung-Hsing University, Taichung, Taiwan, ROC; 1998.
27. Sambrook J, Fritsch EF, Maniatis T, eds. *Molecular cloning: A laboratory manual*. 2nd ed. New York: Cold Spring Harbor Laboratory Press; 1989.
28. Horn U, Strittmatter W, Krebber A, Knupfer U, Kujau M, Wenderoth R, et al. High volumetric yields of functional dimeric miniantibodies in *Escherichia coli*, using an optimized expression vector and high-cell-density fermentation under non-limited growth conditions. *Appl Microbiol Biotechnol* 1996; 46:524-32.
29. Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970;227: 680-5.
30. Hsu WH, Chien FT, Hsu CL, Wang TC, Yuan HS, Wang WC. Expression, crystallization and preliminary X-ray diffraction studies of N-carbamyl-D-amino-acid amidohydrolase from *Agrobacterium radiobacter*. *Acta Cryst D* 1999;55:694-5.

31. Miller GL. Use of dinitrosalicylic acid reagent for determination of reducing sugar. *Anal Chem* 1959;31:426-8.
32. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 1976;72:248-54.
33. Runser S, Ohleyer E. Properties of the hydantoinase from *Agrobacterium* sp. IP I-671. *Biotechnol Lett* 1990;12:259-64.
34. Akesson M, Hagander P, Axelsson JP. Avoiding acetate accumulation in *Escherichia coli* cultures using feedback control of glucose feeding. *Biotechnol Bioeng* 2001;73:223-30.
35. Runser SM, Meyer PC. Purification and biochemical characterization of the hydantoin hydrolyzing enzyme from *Agrobacterium* species: A hydantoinase with no 5,6-dihydropyrimidine amidohydrolase activity. *Eur J Biochem* 1993;213:1315-24.