




Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

ScienceDirect

journal homepage: [www.e-jmii.com](http://www.e-jmii.com)



ORIGINAL ARTICLE

# Fluoroquinolone-resistant and extended-spectrum $\beta$ -lactamase-producing *Escherichia coli* from the milk of cows with clinical mastitis in Southern Taiwan



Yaochi Su <sup>a</sup>, Chang-You Yu <sup>a</sup>, Yilin Tsai <sup>b</sup>, Shao-Hung Wang <sup>b</sup>,  
Chihan Lee <sup>b</sup>, Chishih Chu <sup>b,\*</sup>

<sup>a</sup> Department of Veterinary Sciences, National Chiayi University, Chiayi, Taiwan, ROC

<sup>b</sup> Department of Microbiology, Immunology, and Biopharmaceutics, National Chiayi University, Chiayi, Taiwan, ROC

Received 22 February 2014; received in revised form 1 October 2014; accepted 7 October 2014

Available online 11 November 2014

## KEYWORDS

cow;  
*Escherichia coli*;  
extended-spectrum  
 $\beta$ -lactamase;  
fluoroquinolone;  
mastitis

**Background/Purpose:** *Escherichia coli* is a common pathogen to cause clinical and subclinical mastitis in cows. A total of 57 *E. coli* isolates from raw milk from cows were characterized genetically and biochemically.

**Methods:** Extended-spectrum  $\beta$ -lactamase (ESBL) genes, the mechanism for fluoroquinolone resistance, and variations in virulence genes and genomes of these *E. coli* isolates were investigated by the antimicrobial susceptibility test, simplex and multiplex polymerase chain reaction (PCR), and pulsed-field gel electrophoresis (PFGE).

**Results:** All *E. coli* isolates were resistant to cloxacillin (100%) and to a lesser extent (50%) to tetracycline, neomycin, gentamycin, ampicillin, ceftriaxone, cefotaxime (CTX), and ceftazidime (CAZ). Nearly 70% of the isolates were resistant to at least two antimicrobials and 28.1% carried AmpA and AmpC genes simultaneously. The predominant *bla* gene was *bla*<sub>TEM</sub>, followed by *bla*<sub>CMY</sub>, *bla*<sub>CTX</sub>, *bla*<sub>SHV</sub>, and *bla*<sub>DHA</sub>. Among the six (10.5%) ESBL-producing *E. coli* carrying *bla*<sub>CTX-M15</sub>, *bla*<sub>CTX-M55</sub>, or *bla*<sub>CTX-M14</sub>, two isolates 31 of ST410 in the ST23 complex and 58 of ST167 in the ST10 complex were also resistant to ciprofloxacin, enrofloxacin, and levofloxacin, with mutations at codon 83 from serine to leucine and codon 87 from aspartic acid to asparagine in GyrA and at codon 80 from serine to isoleucine in ParC. These isolates were genetically diverse in pulsotype analysis, lacked toxin genes of human

\* Corresponding author. Department of Microbiology, Immunology, and Biopharmaceutics, National Chiayi University, 300 University Road, Chiayi 60004, Taiwan, ROC.

E-mail address: [cschu@mail.ncyu.edu.tw](mailto:cschu@mail.ncyu.edu.tw) (C. Chu).

pathogenic *E. coli* and carried mostly the prevalent virulence genes *fimH*, *papGII*, and  $\alpha$ -hemolysin.

**Conclusion:** Lacking virulence genes examined, genetic diverse *E. coli* isolates are unrelated to human pathogenic *E. coli*. Enhancing sanitation in milk processing and transportation is needed to eliminate multidrug-resistant (MDR), fluoroquinolone-resistant, and ESBL-producing *E. coli* isolates.

Copyright © 2014, Taiwan Society of Microbiology. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Mastitis in cows and goats affects milk quality, human safety, and the income of the farming industry worldwide. Cases of subclinical mastitis are much higher than those of clinical mastitis. The predominant pathogens causing mastitis are Gram-negative bacteria (25–30%), coagulase-negative *Staphylococcus* (20%), *Staphylococcus aureus* and *Escherichia coli* (5–10% each), *Streptococcus* and *Enterococcus* (2–5%), and other species (< 2%).<sup>1–5</sup> In Taiwan, the predominant pathogens causing mastitis in dairy cows are *E. coli* (16.8%), *Staphylococcus* spp. (23.3%), *Streptococcus* spp. (14.5%), *Enterococcus* spp. (20.1%), and other Gram-negative versus Gram-positive (19.2% vs. 3.8%) pathogens.<sup>6</sup> As normal intestinal flora of the intestine, *E. coli* produces vitamins B and K and inhibits the growth of pathogens in the intestine,<sup>7</sup> and becomes diverse genetic groups and opportunistic pathogens to humans by incorporating different pathogenicity islands with virulence genes, for example, enterotoxigenic *E. coli* (ETEC) with genes for heat-stable toxin and heat-labile toxin,<sup>8</sup> enterohemorrhagic *E. coli* (EHEC) with genes for Shiga-like toxin I and II,<sup>9</sup> and enteropathogenic *E. coli* (EPEC) with genes for bundle-forming pilus.<sup>10</sup> The emergence of multidrug-resistant (MDR) bacteria and the transmission of these bacteria from foods to humans have become serious public health problems in treating their infection.<sup>11,12</sup>

$\beta$ -lactamase can hydrolyze the  $\beta$ -lactam rings of expanded-spectrum cephalosporins (except cephamycins and carbapenems) and monobactams and then inactivates these compounds. Application of third-generation cephalosporins increases extended-spectrum  $\beta$ -lactamase (ESBL) *Enterobacteriaceae*, which are relatively high in food animals and reveal a high genetic diversity,<sup>13,14</sup> indicating that farm animals are a reservoir of ESBL-producing bacteria which can infect humans. Recently, ESBL-producing *E. coli* has been observed in economically important animals in Europe<sup>15,16</sup> and Asia,<sup>17</sup> in meat, fish, and raw milk,<sup>18,19</sup> and in healthy dairy cattle and retail meat in the USA.<sup>20,21</sup> Furthermore, fluoroquinolone- and carbapenem-resistant *E. coli* isolates have been reported in humans. Here, we investigated antimicrobial susceptibility, ESBL-producing genes, mutations in GyrA and ParC for fluoroquinolone resistance, and differences in virulence genes and genomes for mastitis-associated *E. coli* isolates.

## Methods

### Mastitis-associated bacterial species

A total of 412 milk samples from cows with clinical or subclinical mastitis were collected from 13 farms in Yunlin, Chiayi, and Tainan in Southern Taiwan based on the California mastitis test of microcolloid formation<sup>22</sup> and these samples were directly plated on Eosin Methylene Blue (EMB) agar to screen for *E. coli*. Bacterial colonies were further identified as *E. coli* by API 20E and polymerase chain reaction (PCR) amplification of the *E. coli*-specific gene *mdhR*. Hemolysin types of each isolate were determined by blood agar. Informed consent was obtained from all farm owners prior to the start of this study. The use of all bacteria was supervised by the Biological Security Committee of the National Chiayi University, Chiayi, Taiwan in accordance with the laws of Taiwan. The milk samples were collected by county governors based on the guidelines of Animal Use Protocol, the Institutional Animal Care and Use Committee, National Chiayi University.

### Antimicrobial susceptibility test

Antimicrobial susceptibility to ampicillin, cefepime, cefotaxime (CTX), ceftazidime (CAZ), ceftriaxone, ciprofloxacin, cloxacillin, enrofloxacin, ertapenem, gentamicin, levofloxacin, meropenem, neomycin, and tetracycline was determined by the disc diffusion method and the Clinical and Laboratory Standards Institute standard.<sup>23</sup> ESBL isolates were confirmed by determining their resistance to CAZ, CTX, CAZ/clavulanic acid, and CTX/clavulanic acid. *E. coli* ATCC 25922 was used as the reference strain.

### PCR detection of genes for $\beta$ -lactamase, virulence factors, and pathogenic *E. coli*

The primers used to amplify five  $\beta$ -lactamase genes, 12 virulence genes of uropathogenic *E. coli*, and 10 toxin genes of other pathogenic *E. coli* are listed in Table 1.<sup>24–29</sup> Three multiplex PCR sets were developed. Multiplex PCR I was used to detect genes *bla*<sub>SHV</sub>, *bla*<sub>CTX-M3</sub>, *bla*<sub>CTX-M14</sub>, *bla*<sub>TEM</sub>, and *bla*<sub>DHA</sub>. Twelve virulence genes were amplified by multiplex PCR II for *hlyA*, *usp*, *sat*, *fyuA*, *ironN*, *iutA*, and *iucD* and multiplex PCR III for *fimH*,

**Table 1** Primer sequences and PCR product sizes of genes for antimicrobial resistance and virulence factors

PCR Set	Gene	Primer	Nucleotide sequence ( 5' → 3' )	PCR product size (bp)	
Multiplex I <sup>a</sup>	Class A $\beta$ -lactamase				
	<i>bla</i> <sub>CTX-M3</sub>	F	AATCA CTGCG CCAGT TCACG CT	479	
		R	GAACG TTTTCG TCTCC CAGCT GT		
	<i>bla</i> <sub>CTX-M14</sub>	F	TACCG CAGAT AATAC GCAGG TG	355	
		R	CAGCG TAGGT TCAGT GCGAT CC		
	<i>bla</i> <sub>SHV</sub>	F	AACGG AACTG AATGA GGC GC T	141	
		R	TCCAC CATCC ACTGC AGCAG CT		
	<i>bla</i> <sub>TEM</sub>	F	GAAGA TCAGT TGGGT GCACG AGT	520	
		R	CAACT TTATC CGCCT CCATC CAGT		
	Multiplex II	Class C $\beta$ -lactamase			
<i>bla</i> <sub>DHA</sub>		F	AACTT TCACA GGTGT GCTGG GT	405	
		R	CGTAC GCATA CTGGC TTTGC		
Simplex <i>bla</i> <sub>CMY-2</sub>		F	CTGAC AGCCT CTTTC TCCAC A	1100	
		R	CTACG TAGCT GCCAA ATCCA C		
Multiplex II		<i>hlyA</i>	F	AACAA CGATA AGCAC TGTTT TGG	1177
			R	CCATA TAAGC GGTC A TTCCC G	
		<i>fyuA</i>	F	GGCTT TATCC TCTGG CCTT	878
			R	GAAAA CCCAG TCATC GGTGG	
		<i>iroN</i>	F	CTCTG GTGGT GGAAG CC	815
	R		TGTCG GTACA GGCGG TTC		
	<i>usp</i>	F	GGAAA ATGGT CGCTC AGTGG	992	
		R	CTGTA GTGAA TCTCA TCGTG TAGTC		
	<i>sat</i>	F	TCAGA AGCTC AGCGA ATCAT TG	931	
		R	CATTA TCACC AGTAA AACGC ACC		
	<i>iutA</i>	F	CACTC CGGTA CTCCA GTCA	688	
		R	CCTCC AACCA GATGT TCTTC G		
	<i>iucD</i>	F	CCGGA GAAGC CTGAA ATATA TTCA	584	
		R	CCGGA TTGTC ATATG CAGAC C		
Multiplex III	<i>fimH</i>	F	GTTTA TAATT CGAGA ACGGA TAAGC C	494	
		R	GTGCA TAATT TGCCG TTAAT CCC		
	<i>cnf1</i>	F	TTCTT CTGTA CTTC CCCAG	407	
		R	TGAGC GGCAT CTA CTACT ATGAA GT		
	<i>kpsMTII</i>	F	CATCA GACGA TAAGC ATGAG CA	270	
		R	TGCGC ATTTG CTGAT ACTGT		
	<i>papGII</i>	F	GGGCC CCCAA GTAAC TC	188	
		R	GGATG AGCGG GCCTT TG		
	<i>traT</i>	F	CATAA CCACG GTTCA GCCAT C	328	
		R	TTGCA CTGGT CAGTT CCAC		
Specific PCR sets for EHEC <sup>b</sup>	<i>stx1</i>	F	CAGTT AATGT GGTGG CGAAG G	348	
		R	CACCA GACAA TGTA CCGCT G		
	<i>stx2</i>	F	ATCCT ATTCC CGGGA GTTTA CG	584	
		R	GCGTC ATCGT ATACA CAGGA GC		
Specific PCR sets for EPEC <sup>c</sup>	<i>eae</i>	F	TCAAT GCAGT TCCGT TATCA GTT	482	
		R	GTAAA GTCCG TTACC CCAAC CTG		
	<i>bfp</i>	F	GGAAG TCAAA TTCAT GGGGG TAT	300	
		R	GGAAT CAGAC GCAGA CTGGT AGT		
Specific PCR sets for ETEC <sup>c</sup>	<i>Lt</i>	F	GCACA CGGAG CTCCT CAGTC	218	
		R	TCCTT CATCC TTTCA ATGGC TTT		
	<i>stII</i>	F	AAAGG AGAGC TTCGT CACAT TTT	129	
		R	AATGT CCGTC TTGCG TTAGG AC		
Specific PCR sets for EIEC <sup>d</sup>	<i>virF</i>	F	AGCTC AGGCA ATGAA ACTTT GAC	618	
		R	TGGGC TTGAT ATTCC GATAA GTC		
	<i>ipaH</i>	F	CTCGG CACGT TTTAA TAGTC TGG	933	
		R	GTGGA GAGCT GAAGT TTCTC TGC		
Specific PCR set for DAEC <sup>d</sup>	<i>daaE</i>	F	GAACG TTGGT TAATG TGGGG TAA	542	
		R	TATTC ACCGG TCGGT TATCA GT		
Specific PCR set for EAggEC3 <sup>d</sup>	<i>aafII</i>	F	CACAG GCAAC TGA AA TAAGT CTGG	378	
		R	ATTCC CATGA TGCA AGCAC TTC		

Table 1 (continued)

PCR Set	Gene	Primer	Nucleotide sequence ( 5' → 3' )	PCR product size (bp)
Quinolone-resistant determining region	<i>gyrA</i> <sup>e</sup>	F	GCGCGGGCTGTGTTATAATT	519
		R	CCGTGCCGTCATAGTTATCAA	
	<i>parC</i> <sup>f</sup>	F	AAACCTGTTTCAGCGCCGATT	391
		R	GTGGTGCCGTTAAGCAAA	

<sup>a</sup> The PCR method was described by Chia et al.<sup>24</sup> and primers for *bla*<sub>DHA</sub> was designed in this study.

<sup>b</sup> The PCR method was described by Cebula et al.<sup>25</sup>

<sup>c</sup> The PCR method was described by Stacy-Phipps et al.<sup>26</sup>

<sup>d</sup> The method was described by Vidal et al.<sup>27</sup>

<sup>e</sup> The method was described by Cattoir et al.<sup>28</sup>

<sup>f</sup> The method was described by Wada et al.<sup>29</sup>

DAEC = diffuse adherent *Escherichia coli*; EAggEC3 = enteroaggregative *Escherichia coli*; EHEC = enterohemorrhagic *Escherichia coli*; EIEC = enteroinvasive *Escherichia coli*; EPEC = enteropathogenic *Escherichia coli*; ETEC = enterotoxigenic *Escherichia coli*.

*cnf1*, *traT*, *kpsMTII*, and *papGII*. *bla*<sub>CMY-2</sub> was identified by single PCR. A 50- $\mu$ L PCR reagent included 5  $\mu$ L of DNA template, 0.2  $\mu$ M of each primer, 5  $\mu$ L of 10X PCR reaction buffer, 200  $\mu$ M of deoxyribonucleoside triphosphates (dNTPs), and 1.4 U Taq DNA polymerase. The PCR conditions were as follows: predenaturation at 95°C for 5 minutes, 30 cycles of denaturation at 95°C for 1 minute, annealing at 60°C for 1 minute and extension at 72°C for 1 minute, and a final extension at 72°C for 10 minutes. Toxin genes for pathogenic *E. coli* were amplified with previously described PCR conditions.<sup>24–29</sup> All PCR products were separated on 2% agarose gels at 50 V for 120 minutes. After ethidium bromide staining and UV illumination, gel images were recorded and analyzed.

### Sequence analysis of quinolone-resistance determining regions of *gyrA*, *parC*, and *bla*<sub>CTX-M</sub>

The quinolone-resistance determining regions of *gyrA* and *parC* genes were amplified from two fluoroquinolone resistant strains and sensitive strains with the PCR primers listed in Table 1. PCR products were purified and sequenced. The *bla*<sub>CTX-M</sub> was amplified with the primer set in Table 1 and sequenced. The sequences were compared to the NCBI database by the Blast program.

### Pulsed-field gel electrophoresis analysis

The PulseNet standardized laboratory pulsed-field gel electrophoresis (PFGE) protocol for molecular subtyping of *E. coli* O157:H7, non-typhoidal *Salmonella* serotypes, and *Shigella sonnei* was used to analyze the isolates.<sup>30</sup> A total of 10 U of *Xba*I was used for restriction digestion. Digested DNA fragments were separated by the CHEF-DRIII system (BioRad, Hercules, CA, USA) in 1% agarose gel with 0.5X TBE running buffer at 14°C in conditions of 6 V/cm, 120° separation angle, and switching time of 4~70 seconds for 18 hours and another run for 4 hours in the same condition, with only change of voltages to 4 V/cm. PFGE images were analyzed and a unique PFGE pattern was defined as one or two DNA bands differing between PFGE patterns of two isolates.

## Results

### Antimicrobial susceptibility of *E. coli*

*E. coli* from milk samples of the cows with clinical mastitis differed in prevalence of 13.8% (57/412) in total, 51.6% in Yunlin, 10.8% in Chiayi, and 4.1% in Tainan. Based on resistance to 1–12 antimicrobials, the number of antibiograms ranged from 13 in Yunlin to six in Chiayi and Tainan; however, the highest antimicrobial resistance was found in two isolates from Chiayi and Tainan (Table 2). Although all 57 isolates were resistant to cloxacillin, a drug used to treat *S. aureus*, a total of 33.3% of the isolates were susceptible to the remaining 11 antimicrobials, with a prevalence of 64.7% in Chiayi, 37.5% in Tainan, and 15.6% in Yunlin. Prevalence of the resistance to traditional antimicrobials was 51% for tetracycline, 42% for neomycin, and 19% for gentamicin.

Regarding resistance to beta-lactam antimicrobials, 54% isolates were resistant to ampicillin of the penicillin group, while < 25% isolates were resistant to CAZ, CTX, and ceftriaxone of the third-generation cephalosporins, and one isolate was resistant to cefepime (the 4<sup>th</sup>-generation cephalosporin). Furthermore, only six (10.7%) ESBL-producing isolates were identified and all revealed resistance to 7–11 antimicrobials (Table 2). ESBL-producing isolates VMEC31 and VMEC58 were also resistant to fluoroquinolones (ciprofloxacin, enrofloxacin, and levofloxacin) and revealed mutations at codon 83 from serine to leucine and codon 87 from aspartic acid to asparagine in *GyrA*, and at codon 80 from serine to isoleucine in *ParC*. These isolates reduced susceptibility to carbapenems.

### Variations in *bla* genes

Among five *bla* genes encoding AmpA and AmpC lactamase, the prevalent *bla* genes in order were *bla*<sub>TEM</sub> (47%), *bla*<sub>CMY</sub> (40%), *bla*<sub>CTX</sub> [*bla*<sub>CTX-M3-like</sub> (9%) and *bla*<sub>CTX-M14-like</sub> (4%)], *bla*<sub>SHV</sub> (9%), and *bla*<sub>DHA</sub> (2%) (Table 3). Over 70% of the isolates contained at least one *bla* gene, while the number of isolates without the *bla* gene was identical to those resistant only to cloxacillin in Yunlin and Tainan, not Chiayi (Table 3). The predominant *bla* genes were *bla*<sub>TEM</sub> in 43.9%

**Table 2** Prevalence of antimicrobial resistance in different counties

Resistance no.	Resistance (%)											ESBL	Yunlin	Chiayi	Tainan	Total, n (%)		
	Penicillin		Cephalosporin			Fluoroquinolone			GEN	NEO	TET							
	CLO	AMP	CAZ	CTX	CRO	FEP	CIP	ENR									LVX	
1	R	S	S	S	S	S	S	S	S	S	S	S	–	–	5 (15.6)	11 (64.7)	3 (37.5)	19 (33.3)
2	R	S	S	S	S	S	S	S	S	S	S	R	–	–	1 (3.1)	0 (0.0)	0 (0.0)	1 (1.8)
	R	S	S	S	S	S	S	S	S	S	S	R	–	–	2 (6.3)	0 (0.0)	0 (0.0)	2 (3.5)
3	R	R	S	S	S	S	S	S	S	S	S	S	–	–	3 (9.4)	1 (5.9)	0 (0.0)	4 (7.0)
	R	S	S	S	S	S	S	S	S	S	R	R	–	–	2 (6.3)	1 (5.9)	0 (0.0)	3 (5.3)
	R	S	S	S	S	S	S	S	S	R	R	S	–	–	1 (3.1)	0 (0.0)	0 (0.0)	1 (1.8)
	R	R	S	S	S	S	S	S	S	S	S	R	–	–	6 (18.8)	1 (5.9)	0 (0.0)	7 (12.3)
4	R	R	S	S	S	S	S	S	S	S	R	S	–	–	2 (6.3)	0 (0.0)	0 (0.0)	2 (3.5)
	R	R	S	R	S	S	S	S	S	S	R	R	–	–	0 (0.0)	2 (11.8)	0 (0.0)	2 (3.5)
5	R	R	S	R	S	S	S	S	S	S	R	R	–	–	3 (9.4)	0 (0.0)	1 (12.5)	4 (7.0)
	R	R	S	S	S	S	S	S	S	R	R	R	–	–	2 (6.3)	0 (0.0)	1 (12.5)	3 (5.3)
6	R	R	S	R	S	S	S	S	S	R	R	R	–	–	0 (0.0)	0 (0.0)	1 (12.5)	1 (1.8)
	R	R	S	R	R	S	S	S	S	R	R	R	–	–	1 (3.1)	0 (0.0)	0 (0.0)	1 (1.8)
7	R	R	S	R	R	S	S	S	S	R	R	R	+	–	1 (3.1)	0 (0.0)	0 (0.0)	1 (1.8)
8	R	R	R	R	R	S	S	S	S	R	R	R	+	+	3 (9.4)	0 (0.0)	0 (0.0)	3 (5.3)
11	R	R	R	R	R	S	R	R	R	R	R	R	+	+	0 (0.0)	0 (0.0)	1 (12.5)	1 (1.8)
	R	R	R	R	R	R	R	R	R	R	S	R	+	+	0 (0.0)	1 (5.9)	0 (0.0)	1 (1.8)
Total, n (%)	57 (100)	31 (54)	5 (9)	13 (23)	7 (12)	1 (2)	2 (4)	2 (4)	2 (4)	11 (19)	24 (42)	29 (51)	6 (11)	5 (9)	32 (100)	17 (100)	8 (100)	57 (100)

AMP = ampicillin; CAZ = ceftazidime; CLO = cloxacillin; CRO = ceftriaxone; CIP = ciprofloxacin; CTX = cefotaxime; ENR = enrofloxacin; ESBL = extended-spectrum  $\beta$ -lactamase; ETP = ertapenem; FEP = cefepime; GEN = gentamicin; LVX = levofloxacin; NEO = neomycin; TET = tetracycline.

**Table 3** Prevalence of different beta-lactamase genes in three counties

No. of resistant gene	AmpA <i>bla</i>				AmpC <i>bla</i>		ESBL		County			Total, <i>n</i> (%)
	TEM	SHV	CTX-M-3	CTX-M-14	DHA	CMY	CAZ	CTX	Yunlin	Chiayi	Tainan	
0	–	–	–	–	–	–	–	–	5 (15.6)	8 (47.1)	3 (37.5)	16 (28.1)
1	+	–	–	–	–	–	–	–	8 (25.0)	3 (17.6)	2 (25.0)	13 (22.8)
	–	+	–	–	–	–	–	–	1 (3.1)	2 (11.8)	0	3 (5.3)
	–	–	–	+	–	–	–	–	1 (3.1)	0	0	1 (1.8)
	–	–	–	–	+	–	–	–	1 (3.1)	0	0	1 (1.8)
	–	–	–	–	–	+	–	–	4 (12.5)	3 (17.6)	0	7 (12.3)
2	+	–	–	–	–	+	–	–	7 (21.9)	0	1 (12.5)	8 (14.0)
	–	–	+	–	–	+	+	+	0	1 (5.9)	0	1 (1.8)
	–	–	–	+	–	+	–	+	1 (3.1)	0	0	1 (1.8)
3	+	+	–	–	–	+	–	–	1 (3.1)	0	1 (12.5)	2 (3.5)
	+	–	+	–	–	+	+	+	3 (9.4)	0	1 (12.5)	4 (7.0)
Total, <i>n</i> (%)	27 (47)	5 (9)	5 (9)	2 (4)	1 (2)	23 (40)	5 (9)	6 (11)	32	17	8	57 (100)

CAZ = ceftazidime; CMY = ; CTX = cefotaxime; DHA = ; ESBL = extended-spectrum  $\beta$ -lactamase; SHV = ; TEM = .

of isolates carrying one *bla* gene, *bla*<sub>TEM</sub> and *bla*<sub>CMY</sub> in 17.6% of isolates with two *bla* genes, and *bla*<sub>TEM</sub>, *bla*<sub>CTX-M3</sub>, and *bla*<sub>CMY</sub> in 10.5% of isolates with three *bla* genes. The latter two groups all contained AmpC *bla*<sub>CMY</sub> and at least one AmpA *bla* gene. Isolates with three *bla* genes were only observed in Yunlin and Tainan. In contrast to CAZ<sup>ESBL</sup> and CTX<sup>ESBL</sup> phenotypes of *E. coli* carrying *bla*<sub>CTX-M3-like</sub>, *E. coli* with *bla*<sub>CTX-M14-like</sub> exhibited only a CTX<sup>ESBL</sup> phenotype or non-ESBL phenotype. Sequence analysis of *bla*<sub>CTX-M3-like</sub> genes determined *bla*<sub>CTX-M55</sub> for isolates 39, 42, 43, and 58 and *bla*<sub>CTX-M15</sub> for isolate 31.

### Differences in virulence genes

Examination of hemolysin types found  $\alpha$ -hemolysin in 96.7% (55/57) isolates and  $\beta$ - and  $\gamma$ -hemolysin in one isolate, respectively. PCR amplification of toxin genes only identified *eae* of EPEC in one isolate and lacked other toxin genes of EHEC, ETEC, enteroinvasive *E. coli* (EIEC), diffuse adherent *E. coli* (DAEC), and enteroaggregative *E. coli* (EAggEC). Among the other 13 virulence genes for urinary tract infection (UTI)-associated *E. coli* in humans, almost all isolates carried *fimH* encoding type 1 fimbriae and *papGII* encoding P fimbriae, and < 10% of isolates consisted of the *kpsMTII* encoding capsule and *cnf1* and *hlyA* encoding cytotoxin (Table 4). Both capsule and cytotoxin genes were only present in the isolates from Yunlin (14%, 8/57) and Chiayi (2%, 1/57) and were absent in Tainan, while both fimbriae and cytotoxin genes were only observed in the isolates from Yunlin (7%, 4/57) and Tainan (2%, 1/57). Furthermore, siderophore genes were present in < 40% of isolates and ranged from 10.5% (6/57) for *iroN*, encoding the aerobactin receptor, 40.3% (23/57) for *iutA* and 15.8% for *iucD*, encoding salmochelin, and 17.5% for *fyuA*, encoding the yersiniabactin receptor. However, almost all isolates with *iucD* carried *iutA*.

### Phylogenetic analysis

PFGE analysis separated 57 isolates into 39 pulsotypes and nontypeable (Fig. 1 and Table 3). Clonal dissemination was

only observed in pulsotypes 5, 16, 22, and nontypeable in Yunlin, and in pulsotypes 20, 25, 32, and nontypeable in Chiayi. These nontypeable isolates differed in their virulence genes. Pulsotypes of two fluoroquinolone-resistant strains were pulsotype 28 for isolate 31 and nontypeable for isolate 58.

### Discussion

*E. coli* are important pathogenic sources for cow mastitis with a prevalence of 5–10% in South Eastern Ethiopia,<sup>1</sup> 15.9% for clinical mastitis in Estonia,<sup>31</sup> and 4.8% for sub-clinical mastitis.<sup>32</sup> In this study, genetically diverse *E. coli* isolates were the predominant pathogens from the milk of cows with clinical mastitis (Fig. 1), suggesting that these *E. coli* isolates may originate from the environment. Virulence factors, such as S and P fimbriae and CNF1, are important for the severity and persistence of mastitis.<sup>33</sup> Examining the virulence and toxin genes for disease in cattle, these *E. coli* isolates carried *fimH*, *papGII*, and  $\alpha$ -hemolysin genes more prevalent than other virulence genes (> 80% vs. < 40%; Fig. 1, Table 3), implying that these three genes are important for clinical mastitis in cows and that these *E. coli* isolates was not associated with EPEC, EHEC, ETEC, EIEC, DAEC, and EAggEC pathogenic to humans.


Subtherapeutic antimicrobials are frequently used in animal feed, leading to an increase in antimicrobial-resistant bacteria in the animal. In feedlot cattle, given feed with subtherapeutic antimicrobials, the isolates increased resistance to sulfamethoxazole and chloramphenicol, and did not change other antimicrobial resistance patterns.<sup>34</sup> Recently, MDR *E. coli* isolates have emerged from cows with mastitis. Antimicrobial resistance of mastitis-associated *E. coli* isolates differs among countries from low resistance in Sweden and higher resistance in France.<sup>35</sup> Additionally, *E. coli* isolates from the feedlot cattle were less resistant to ampicillin (5.9–24.3%), kanamycin (15.9%), streptomycin (5.9–15.6%), tetracycline (5.9–13.5%), susceptible to cefotaxime, ceftiofur, chloramphenicol, gentamicin, and nalidixic acid, and not ESBL producers,<sup>32,34</sup> while other research reported the isolates

**Table 4** Prevalence of virulence genes in three counties

No. factor	Fimbriae		Capsule	Cytotoxin		Siderophore			Pulsotype	Number	County			Sum
	<i>fimH</i> + <i>papGII</i>	<i>kpsMTII</i>	<i>cnf1</i> + <i>hlyA</i>	<i>iroN</i>	<i>iutA</i>	<i>iucD</i>	<i>fyuA</i>	Yunlin			Chiayi	Tainan		
0	-	-	-	-	-	-	-	-	18;24/25/ 26/31;40	1;1/2/ 1/1/1;1	1	5	1	7
2	-	-	+	-	-	-	-	-	1/22	1/2	3			17
	-	-	-	-	-	-	-	-	4/7/8/11/ 19/23;20/ 32/33;38	1/1/1/1/ 1/1/1/1; 2/3/1;1	6	6	1	
3	+	-	-	-	+	+	-	-	ND	1				1
	+	-	-	-	+	-	-	-	8/14/16/ 17/21,29/ ND;37	1/1/2/ 1/1;1/ 3;1	6	4	1	19
	+	-	-	-	-	-	+	-	3/ND;34	1/2;1	3			1
4	+	-	-	-	-	-	-	-	27;36	1;1		1		1
	+	+	-	-	-	-	-	-	10/15	1/1	2			
	+	-	+	-	-	-	-	-	20	1	1			3
5	+	-	-	-	+	+	-	-	35	1			1	
	+	+	-	-	+	-	+	-	3	1	1			8
6	+	+	-	-	+	+	-	-	6	1	1			
	+	-	-	-	+	+	+	-	13	1	1			
	+	-	+	+	-	+	-	-	12	1	1			
7	+	+	-	-	+	+	-	-	2/5	1/3	4			
	+	-	-	+	+	+	+	-	39	1			1	3
	+	-	+	+	-	-	+	-	9	1	1			
7	+	+	-	+	+	+	+	9	1	1				
Total	48 (84.2)	9 (15.8)	8 (14.0)	6 (10.5)	23 (40.3)	9 (15.8)	10 (17.5)		28	1	32	17	8	57

more resistant to ampicillin (100%), chloramphenicol (75.4%), gentamycin (54%), neomycin (43.2%), streptomycin (97.3%), sulfonamide (98.6%), and tetracycline (100%).<sup>36</sup> However, resistance to fluoroquinolones or cephalosporins is still uncommon among coliforms isolated from bovine mastitis. Although < 50% isolates were resistant to ampicillin, gentamycin, neomycin, and tetracycline, an ESBL producer was identified in this study (Table 3).

Beta-lactam and fluoroquinolone antimicrobials have been frequently used to treat bacterial infections in veterinary hospitals and farms. The use of a third-generation cephalosporin, ceftiofur, in dairy cows and cattle reduces the susceptibility to ceftriaxone in *E. coli* and does not have an effect on the maintenance and dissemination of *bla*<sub>CMY-2</sub>-bearing plasmid.<sup>37,38</sup> However, the application of these cephalosporins may increase ESBL-producing *E. coli*



**Figure 1.** Pulsed-field gel electrophoresis analysis of clinical *Escherichia coli* isolates associated with cow mastitis.

**Table 5** Characterization of *Escherichia coli* isolates from cattle worldwide

Country	Animal condition	Source	CTX-M type	Other <i>bla</i> genes	ESBL prevalence (%)	Refs
China	Healthy	Rectal	CTX-M14, CTX-M55			44
Switzerland	Healthy	Feces, Beef Milk	CTX-M1, 14,15, 117 CTX-M1	<i>bla</i> <sub>TEM</sub> <i>bla</i> <sub>TEM</sub>	13.7	14
France	Healthy	Feces	CTX-M9 group CTX-M1	<i>bla</i> <sub>TEM</sub> <i>bla</i> <sub>TEM-71</sub>	17.1	45 40
		Soil, feces				
	Diarrhea, septicemia	Feces, Blood	CTX-M1 group (65.7%), CTX-M2 group (3.9%), CTX-M9 group (27%)	<i>bla</i> <sub>TEM</sub> (3.4%)	100	46
Japan	Diseased		CTX-M2	<i>bla</i> <sub>TEM</sub> , <i>bla</i> <sub>CMY-2</sub>	2.8	43
	Healthy	Rectal	CTX-M		12.5	47
Korea	Healthy	Feces	CTX-M		0.2	48
		Feces, Milk	CTX-M14, CTX-M15, CTX-M32	<i>bla</i> <sub>TEM-1</sub>		49
Ireland	Diseased		CTX-M2	<i>bla</i> <sub>CMY-2</sub>		36
United Kingdom	Healthy	Feces	CTX-M1, 14, 15			39
		Milk	CTX-M15	<i>bla</i> <sub>TEM-1</sub>		50
Taiwan	Clinical mastitis	Milk	CTX-M14, 15, 55	<i>bla</i> <sub>CMY</sub> , <i>bla</i> <sub>TEM</sub> , <i>bla</i> <sub>SHV</sub>	11	This study

ESBL = extended-spectrum  $\beta$ -lactamase.

from animals, such as cattle, worldwide<sup>14,39–43</sup> (Tables 4 and 5)<sup>14,36,39,40,43–50</sup>. The ESBL-producing isolates were more prevalent in diseased cattle than healthy cattle and were obtained from different sources. The *bla*<sub>CTX</sub> is the most important ESBL-related gene for ESBL-producing *E. coli* from cattle and its genotypes are associated with the geographic area (Tables 3 and 5). The prevalent genotypes of the *bla*<sub>CTX</sub> are *bla*<sub>CTX-M14</sub> and *bla*<sub>CTX-M15</sub> worldwide. However, in this study, *bla*<sub>CTX-M55</sub> in the isolates from Yunlin and Tainan was the most prevalent genotype and has been reported from cattle in China. The *bla*<sub>CTX-M55</sub> isolates were genetically identical from the same farm in Yunlin, implying clonal dissemination of *bla*<sub>CTX-M55</sub> *E. coli* in Yunlin.

In the present study, we determined *bla*<sub>CTM-M14</sub>, *bla*<sub>CTM-M15</sub>, and *bla*<sub>CTM-M55</sub> responsible for ESBL in *E. coli*. However, *bla*<sub>CTM-M14</sub> could not present simultaneously with *bla*<sub>CTM-M15</sub> or *bla*<sub>CTM-M55</sub> in the same isolate, indicating that *bla*<sub>CTM-M14</sub> and *bla*<sub>CTM-M15</sub> or *bla*<sub>CTM-M55</sub> may exist mutually exclusively, possibly on different incompatibility plasmids, and therefore, the transferability of *bla*<sub>CTM-M14</sub> and *bla*<sub>CTM-M15</sub> or *bla*<sub>CTM-M55</sub> associated elements need to be confirmed. Additionally, some MDR ESBL-producing isolates carried AmpA and AmpC *bla* genes simultaneously, such as *bla*<sub>CTX-M2</sub> and *bla*<sub>CMY-2</sub>, in hospitalized animals (Table 5).<sup>35</sup> In the present study, 28.1% isolates carried AmpA and AmpC genes simultaneously (Table 3). Carbapenem-resistant *E. coli* can occur in ESBL- and AmpC-producing *E. coli* isolates with mutations in *ompF* and *ompC* genes,<sup>51,52</sup> or be through introduction of the chromosomal- and plasmid-encoded carbapenamase, to become carbapenem-resistant.<sup>53</sup>

MDR fluoroquinolone-resistant ESBL isolates 31 and 58 with both AmpA and AmpC genes, differed in *bla*<sub>CTX-M</sub> genotype and pulsotypes and were located in Chiayi and Tainan, respectively. These results demonstrate different origins of these two isolates. The presence of these isolates

in current cow farms and the mechanism of reduced susceptibility to carbapenem need to be further investigated. To avoid the emergence of such resistant isolates and to prevent the spreading of these isolates, the use of cephalosporins and fluoroquinolones should be limited in treating cow infection isolates, and enhanced sanitation of milk processing and transportation is needed.

In conclusion, genetically divergent *E. coli* isolates were less resistant to traditional antimicrobials and not related to human pathogens. Emergence of MDR fluoroquinolone-resistant ESBL producers with reduced susceptibility to carbapenem may threaten the public health.

## Conflicts of interest

All authors declare that they have no conflicts of interest relevant to this article.

## Acknowledgments

We would like to acknowledge grants from the Council of Agriculture (99AS-9.6.1-BQ-B3), and the National Science Council (100-2320-B-415-002), Executive Yuan of the Republic of China, Taiwan.

## References

- Lakew M, Tolosa T, Tigre W. Prevalence and major bacterial causes of bovine mastitis in Asella, South Eastern Ethiopia. *Trop Anim Health Prod* 2009;41:525–30.
- Quinn AK, Vermunt JJ, Twiss DP. *Arcanobacterium pyogenes* mastitis in a 18-month-old heifer. *NZ Vet J* 2002;50:167–8.
- Apparao MD, Ruegg PL, Lago A, Godden S, Bey R, Leslie K. Relationship between in vitro susceptibility test results and



- treatment outcomes for gram-positive mastitis pathogens following treatment with cephapirin sodium. *J Dairy Sci* 2009; **92**:2589–97.
4. Apparao D, Oliveira L, Ruegg PL. Relationship between results of in vitro susceptibility tests and outcomes following treatment with pirlimycin hydrochloride in cows with subclinical mastitis associated with gram-positive pathogens. *J Am Vet Med Assoc* 2009; **234**:1437–46.
  5. Shum LWC, McConnel CS, Gunna AA, Housea JK. Environmental mastitis in intensive high-producing dairy herds in New South Wales. *Aust Vet J* 2009; **87**:469–75.
  6. Ma YP, Chang SK, Chou CC. Characterization of bacterial susceptibility isolates in sixteen dairy farms in Taiwan. *J Dairy Sci* 2006; **89**:4573–82.
  7. Hudault S, Guignot J, Servin A. *Escherichia coli* strains colonising the gastrointestinal tract protect germfree mice against *Salmonella typhimurium* infection. *Gut* 2001; **49**:47–55.
  8. You J, Xu Y, He M, McAllister TA, Thacker PA, Li X, et al. Protection of mice against enterotoxigenic *E. coli* by immunization with a polyvalent enterotoxin comprising a combination of LTb, STa, and STb. *Appl Microbiol Biotechnol* 2011; **89**:1885–93.
  9. Golan L, Gonen E, Yagel S, Rosenshine I, Shpigel NY. Enterohemorrhagic *Escherichia coli* induce attaching and effacing lesions and hemorrhagic colitis in human and bovine intestinal xenograft models. *Lett Appl Microbiol* 2010; **51**:499–503.
  10. Bugarel M, Martin A, Fach P, Beutin L. Virulence gene profiling of enterohemorrhagic and enteropathogenic *Escherichia coli* strains: a basis for molecular risk assessment of typical and atypical EPEC strains. *BMC Microbiol* 2011; **11**:142.
  11. Angulo FJ, Nargund VN, Chiller TC. Evidence of an association between use of antimicrobial agents in food animals and antimicrobial resistance among bacteria isolated from humans and the human health consequences of such resistance. *J Vet Med B* 2004; **51**:374–9.
  12. Phillips I, Casewell M, Cox T, De GB, Friis C, Jones R, et al. Does the use of antibiotics in food animals pose a risk to human health? A critical review of published data. *J Antimicrob Chemother* 2004; **53**:28–52.
  13. Blanc V, Mesa R, Saco M, Lavilla S, Prats G, Miro E, et al. ESBL- and plasmidic class C  $\beta$ -lactamase producing *E. coli* strains isolated from poultry, pig and rabbit farms. *Vet Microbiol* 2006; **118**:299–304.
  14. Geser N, Stephan R, Hächler H. Occurrence and characteristics of extended spectrum  $\beta$ -lactamase (ESBL) producing *Enterobacteriaceae* in food producing animals, minced meat and raw milk. *BMC Vet Res* 2012; **8**:21.
  15. Cortés P, Blanc V, Mora A, Dahbi G, Blanco JE, Blanco M, et al. Isolation and characterization of potentially pathogenic antimicrobial-resistant *Escherichia coli* strains from chicken and pig farms in Spain. *Appl Environ Microbiol* 2010; **76**:2799–805.
  16. Gonçalves A, Torres C, Silva N, Carneiro C, Radhouani H, Coelho C, et al. Genetic characterization of extended-spectrum  $\beta$ -lactamases in *Escherichia coli* isolates of pigs from a Portuguese intensive swine farm. *Foodborne Pathog Dis* 2010; **7**:1569–73.
  17. Tian GB, Wang HN, Zou LK, Tang JN, Zhao YW, Ye MY, et al. Detection of CTX-M-15, CTX-M-22, and SHV-2 extended-spectrum  $\beta$ -lactamases (ESBLs) in *Escherichia coli* fecal sample isolates from pig farms in China. *Foodborne Pathog Dis* 2009; **6**:297–304.
  18. Hammad AM, Ahmed AM, Ishida Y, Shimamoto T. First characterization and emergence of SHV-60 in raw milk of a healthy cow in Japan. *J Vet Med Sci* 2008; **70**:1269–72.
  19. Jensen LB, Hasman H, Agersø Y, Emborg HD, Aarestrup FM. First description of an oxyimino-cephalosporin-resistant, ESBL-carrying *Escherichia coli* isolated from meat sold in Denmark. *J Antimicrob Chemother* 2006; **57**:793–4.
  20. Doi Y, Paterson DL, Egea P, Pascual A, Lopez-Cerero L, Navarro MD, et al. Extended-spectrum and CMY-type beta-lactamase-producing *Escherichia coli* in clinical samples and retail meat from Pittsburgh, USA and Seville, Spain. *Clin Microbiol Infect* 2010; **16**:33–8.
  21. Wittum TE, Mollenkopf DF, Daniels JB, Parkinson AE, Mathews JL, Fry PR, et al. CTX-M-type extended-spectrum  $\beta$ -lactamases present in *Escherichia coli* from the feces of cattle in Ohio, United States. *Foodborne Pathog Dis* 2010; **7**:1575–9.
  22. Schalm OW, Noorlander DO. Experiments and observations leading to the development of the California Mastitis Test. *J Am Vet Med Assoc* 1957; **130**:199.
  23. Clinical and Laboratory Standards Institute. *Performance standards for antimicrobial susceptibility tests; approved standard*. 12th ed. 2010. M100-S20-U. Wayne, Pennsylvania, USA.
  24. Chia JH, Chu C, Su LH, Chiu CH, Kuo AJ, Sun CF, Wu TL. Development of a multiplex PCR and SHV melting-curve mutation detection system for detection of some SHV and CTX-M beta-lactamases of *Escherichia coli*, *Klebsiella pneumoniae*, and *Enterobacter cloacae* in Taiwan. *J Clin Microbiol* 2005; **43**:4486–91.
  25. Cebula TA, Payne WL, Feng P. Simultaneous identification of strains of *Escherichia coli* serotype O157:H7 and their Shiga-like toxin type by mismatch amplification mutation assay-multiplex PCR. *J Clin Microbiol* 1995; **33**:248–50.
  26. Stacy-Phipps S, Mecca JJ, Weiss JB. Multiplex PCR assay and simple preparation method for stool specimens detect enterotoxigenic *Escherichia coli* DNA during the course of infection. *J Clin Microbiol* 1995; **33**:1054–9.
  27. Vidal M, Kruger E, Durán C, Lagos R, Levine M, Prado V, et al. Single multiplex PCR assay to identify simultaneously the six categories of diarrheagenic *Escherichia coli* associated with enteric infections. *J Clin Microbiol* 2005; **43**:5362–5.
  28. Cattoir V, Lesprit P, Lascols C, Denamur E, Legrand P, Soussy CJ, et al. *In vivo* selection during ofloxacin therapy of *Escherichia coli* with combined topoisomerase mutations that confer high resistance to ofloxacin but susceptibility to nalidixic acid. *J Antimicrob Chemother* 2006; **58**:1054–7.
  29. Wada K, Kariyama R, Mitsuahara R, Uehara S, Watanabe T, Monden K, et al. Experimental and clinical studies on fluoroquinolone-insusceptible *Escherichia coli* isolated from patients with urinary tract infections from 1994 to 2007. *Acta Med Okayama* 2009; **63**:263–72.
  30. Ribot EM, Fair MA, Gautom R, Cameron DN, Hunter SB, Swaminathan B, et al. Standardization of pulsed-field gel electrophoresis protocols for the subtyping of *Escherichia coli* O157:H7, *Salmonella*, and *Shigella* for PulseNet. *Foodborne Pathog Dis* 2006; **3**:59–67.
  31. Persson Y, Nyman AK, Grönlund-Andersson U. Etiology and antimicrobial susceptibility of udder pathogens from cases of subclinical mastitis in dairy cows in Sweden. *Acta Vet Scand* 2011; **53**:36.
  32. Kalmus P, Aasmäe B, Kärssin A, Orro T, Kask K. Udder pathogens and their resistance to antimicrobial agents in dairy cows in Estonia. *Acta Vet Scand* 2011; **53**:4.
  33. Lehtolainen T, Pohjanvirta T, Pyörälä S, Pelkonen S. Association between virulence factors and clinical course of *Escherichia coli* mastitis. *Acta Vet Scand* 2003; **44**:203–5.
  34. Mirzaagha P, Louie M, Sharma R, Yanke LJ, Topp E, McAllister TA. Distribution and characterization of ampicillin- and tetracycline-resistant *Escherichia coli* from feedlot cattle fed subtherapeutic antimicrobials. *BMC Microbiol* 2011; **11**:78.
  35. Suojala L, Kaartinen L, Pyörälä S. Treatment for bovine *Escherichia coli* mastitis – an evidence-based approach. *J Vet Pharmacol Therap* 2013; **36**:521–31.

36. Karczmarczyk M, Abbott Y, Walsh C, Leonard N, Fanning S. Characterization of multidrug-resistant *Escherichia coli* isolates from animals presenting at a university veterinary hospital. *Appl Environ Microbiol* 2011;**77**:7104–12.
37. Daniels JB, Call DR, Hancock D, Sischo WM, Baker K, Besser TE. Role of ceftiofur in selection and dissemination of *bla*<sub>CMY-2</sub>-mediated cephalosporin resistance in *Salmonella enterica* and commensal *Escherichia coli* isolates from cattle. *Appl Environ Microbiol* 2009;**75**:3648–55.
38. Tragesser LA, Wittum TE, Funk JA, Winokur PL, Rajala-Schultz PJ. Association between ceftiofur use and isolation of *Escherichia coli* with reduced susceptibility to ceftriaxone from fecal samples of dairy cows. *Am J Vet Res* 2006;**67**:1696–700.
39. Horton RA, Randall LP, Snary EL, Cockrem H, Lotz S, Wearing H, et al. Fecal carriage and shedding density of CTX-M extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* in cattle, chickens, and pigs: implications for environmental contamination and food production. *Appl Environ Microbiol* 2011;**77**:3715–9.
40. Hartmann A, Locatelli A, Amoureux L, Depret G, Jolivet C, Gueneau E, et al. Occurrence of CTX-M producing *Escherichia coli* in soils, cattle, and farm environment in France (Burgundy Region). *Front Microbiol* 2012;**3**:83.
41. Guerra B, Avsaroglu MD, Junker E, Schroeter A. Detection and characterisation of ESBLs in German *Escherichia coli*, isolated from animal, foods, and human origin between 2001–2006. *Int J Antimicrob Agents* 2007;**S270**–1.
42. Briñas L, Moreno MA, Teshager T, Sáenz Y, Porrero MC, Domínguez L, et al. Monitoring and characterization of extended-spectrum beta-lactamases in *Escherichia coli* strains from healthy and sick animals in Spain in 2003. *Antimicrob Agents Chemother* 2005;**49**:1262–4.
43. Asai T, Masani K, Sato C, Hiki M, Usui M, Baba K, et al. Phylogenetic groups and cephalosporin resistance genes of *Escherichia coli* from diseased food-producing animals in Japan. *Acta Vet Scand* 2011;**53**:52.
44. Zheng H, Zeng Z, Chen S, Liu Y, Yao Q, Deng Y, et al. Prevalence and characterisation of CTX-M  $\beta$ -lactamases amongst *Escherichia coli* isolates from healthy food animals in China. *Int J Antimicrob Agents* 2012;**39**:305–10.
45. Geser N, Stephan R, Kuhnert P, Zbinden R, Kaeppli U, Cernela N, et al. Fecal carriage of extended-spectrum  $\beta$ -lactamase-producing *Enterobacteriaceae* in swine and cattle at slaughter in Switzerland. *J Food Prot* 2011;**74**:446–9.
46. Valat C, Auvray F, Forest K, Métayer V, Gay E, de Garam CP, et al. Phylogenetic grouping and virulence potential of extended-spectrum- $\beta$ -lactamase-producing *Escherichia coli* strains in cattle. *Appl Environ Microbiol* 2012;**78**:4677–82.
47. Hiroi M, Yamazaki F, Harada T, Takahashi N, Iida N, Noda Y, et al. Prevalence of extended-spectrum  $\beta$ -lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* in food-producing animals. *J Vet Med Sci* 2012;**74**:189–95.
48. Tamang MD, Nam HM, Kim SR, Chae MH, Jang GC, Jung SC, et al. Prevalence and molecular characterization of CTX-M  $\beta$ -lactamase-producing *Escherichia coli* isolated from healthy swine and cattle. *Foodborne Pathog Dis* 2013;**10**:13–20.
49. Tamang MD, Nam HM, Gurung M, Jang GC, Kim SR, Jung SC, et al. Molecular characterization of CTX-M  $\beta$ -lactamase and associated addiction systems in *Escherichia coli* circulating among cattle, farm workers, and the farm environment. *Appl Environ Microbiol* 2013;**79**:3898–905.
50. Timofte D, Maciuca IE, Evans NJ, Williams H, Wattret A, Fick JC, et al. Detection and molecular characterisation of *Escherichia coli* CTX-M-15 and *Klebsiella pneumoniae* SHV-12 producing  $\beta$ -lactamases isolated from cases of bovine mastitis in the United Kingdom. *Antimicrob Agents Chemother* 2014;**58**:789–94.
51. Oteo J, Delgado-Iribarren A, Vega D, Bautista V, Rodríguez MC, Velasco M, et al. Emergence of imipenem resistance in clinical *Escherichia coli* during therapy. *Int J Antimicrob Agents* 2008;**32**:534–7.
52. Chia JH, Siu LK, Su LH, Lin HS, Kuo AJ, Lee MH, et al. Emergence of carbapenem-resistant *Escherichia coli* in Taiwan: resistance due to combined CMY-2 production and porin deficiency. *J Chemother* 2009;**21**:621–6.
53. Queenan AM, Bush K. Carbapenemases: the versatile  $\beta$ -lactamases. *Clin Microbiol Rev* 2007;**20**:440–58.